

RB Health (US) LLC

-PE Product Labels –

-continued

MAXIMUM STRENGTH* NDC 63824-536-66

060616 3036245

Mucinex[®]

FAST-MAX[®]

CONGESTION & HEADACHE

Acetaminophen – Pain Reliever/Fever Reducer
Guaifenesin – Expectorant
Phenylephrine HCl – Nasal Decongestant

- ✓ **Relieves Headache & Fever**
- ✓ **Relieves Nasal & Chest Congestion**
- ✓ **Thins & Loosens Mucus**

6 FL OZ (180mL) **FOR AGES 12+**

Drug Facts

Active ingredients (in each 20 mL)

Purposes

Acetaminophen 650 mg.....	Pain reliever/ fever reducer
Guaifenesin 400 mg.....	Expectorant
Phenylephrine HCl 10 mg....	Nasal decongestant

Uses

- temporarily relieves these common cold and flu symptoms:
 - nasal congestion
 - headache
 - minor aches and pains
 - sinus congestion and pressure
- temporarily reduces fever
- temporarily promotes nasal and/or sinus drainage
- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 6 doses in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen

Drug Facts (continued)

- 3 or more alcoholic drinks daily while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
 - blisters
 - rash
- If a skin reaction occurs, stop use and seek medical help right away.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- liver disease
- heart disease
- diabetes
- high blood pressure
- thyroid disease
- trouble urinating due to an enlarged prostate gland
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema

PEEL HERE

Drug Facts (continued)

- cough that occurs with too much phlegm (mucus)

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin

When using this product do not use more than directed

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back, or occurs with fever, rash, or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- do not take more than directed (see Overdose warning)

Drug Facts (continued)

- do not take more than 6 doses in any 24-hour period
- measure only with dosing cup provided
- do not use dosing cup with other products
- dose as follows or as directed by a doctor
- adults and children 12 years of age and over: 20 mL in dosing cup provided every 4 hours
- children under 12 years of age: do not use

Other information

- each 20 mL contains: sodium 12 mg
- store between 20-25°C (68-77°F)
- do not refrigerate

Inactive ingredients anhydrous citric acid, edetate disodium, FD&C blue no. 1, FD&C red no. 40, flavors, glycerin, propyl gallate, propylene glycol, purified water, sodium benzoate, sorbitol, sucralose, trisodium citrate dihydrate[†], xanthan gum[†] may contain this ingredient

Questions?

1-866-MUCINEX (1-866-682-4639)
You may also report side effects to this phone number.



HEALTH • HYGIENE • HOME

Please visit our website
www.mucinex.com

**PEEL CORNER TO READ COMPLETE
DRUG FACTS AND INFORMATION**

***Per 4-hour dose**

**Tamper evident: Do not use if neckband
on bottle cap is broken or missing.**



Dist. by: Reckitt Benckiser
Parsippany, NJ 07054-0224
©2018 RB

011018

LOT:
EXP:
MADE IN:
3055005



Drug Facts		Drug Facts (continued)	
Active Ingredients (in each liquid gel)	Purposes	<ul style="list-style-type: none"> ■ persistent or chronic cough such as occurs with smoking, asthma, or emphysema ■ cough that occurs with too much phlegm (mucus) 	
Acetaminophen 325 mg.....	Pain reliever/fever reducer	Ask a doctor or pharmacist before use if you are	

Dextromethorphan HBr 10 mg.....Cough suppressant Phenylephrine HCl 5 mg.....Nasal decongestant	
Uses ■ temporarily relieves these common cold and flu symptoms: ■ nasal congestion ■ cough ■ headache ■ minor aches and pains ■ sinus congestion and pressure ■ temporarily reduces fever ■ temporarily promotes nasal and/or sinus drainage	
Warnings Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take: ■ more than 12 liquid gels in 24 hours, which is the maximum daily amount ■ with other drugs containing acetaminophen ■ 3 or more alcoholic drinks daily while using this product Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin reddening ■ blisters ■ rash If a skin reaction occurs, stop use and seek medical help right away.	
Do not use ■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. ■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.	
Ask a doctor before use if you have ■ liver disease ■ heart disease ■ diabetes ■ high blood pressure ■ thyroid disease ■ trouble urinating due to an enlarged prostate gland ►	
taking the blood thinning drug warfarin When using this product do not use more than directed Stop use and ask a doctor if ■ nervousness, dizziness, or sleeplessness occur ■ pain, nasal congestion, or cough gets worse or lasts more than 7 days ■ fever gets worse or lasts more than 3 days ■ redness or swelling is present ■ new symptoms occur ■ cough comes back, or occurs with rash or headache that lasts. These could be signs of a serious condition.	
If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.	
Directions ■ do not take more than directed (see Overdose warning) ■ do not take more than 12 liquid gels in any 24-hour period ■ adults and children 12 years of age and older: take 2 liquid gels every 4 hours ■ children under 12 years of age: do not use	
Other information ■ store at 20-25°C (68-77°F) ■ avoid excessive heat	
Inactive Ingredients FD&C red no. 40, FD&C yellow no. 6, gelatin, glycerin, lecithin (soy), mineral oil, polyethylene glycol, povidone, propylene glycol, shellac, sorbitol sorbitan solution, titanium dioxide, water	
Questions? 1-866-MUCINEX (1-866-682-4639) You may also report side effects to this phone number.	



2987

Questions?

1-866-MUCINEX (1-866-682-4639)

You may also report side effects
to this phone number.

LIFT HERE TO OPEN



**Tamper evident: Do not use if outer package
is opened or if blister is torn or broken.**

Drug Facts**Active ingredients (in each caplet) Purposes**

Acetaminophen 325 mg... Pain reliever/fever reducer
Guaifenesin 200 mg.....Expectorant
Phenylephrine HCl 5 mg.....Nasal decongestant

Uses

- temporarily relieves these common cold and flu symptoms:
 - nasal congestion
 - fever
 - headache
 - minor aches and pains
 - sinus congestion and pressure
- temporarily promotes nasal and/or sinus drainage
- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 12 caplets in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks daily while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
 - blisters
 - rash
- If a skin reaction occurs, stop use and seek medical help right away.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- liver disease
- heart disease
- diabetes
- high blood pressure
- thyroid disease

Drug Facts (continued)

- trouble urinating due to an enlarged prostate gland
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- cough that occurs with too much phlegm (mucus)

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin

When using this product do not use more than directed

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back, or occurs with fever, rash, or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- **do not take more than directed (see Overdose warning)**
- do not take more than 12 caplets in any 24-hour period
- adults and children 12 years of age and over: take 2 caplets every 4 hours
- children under 12 years of age: do not use

Other information

- store between 20-25°C (68-77°F)

Inactive ingredients corn starch, croscopovidone, FD&C red no. 40 aluminum lake, FD&C yellow no. 6 aluminum lake, magnesium stearate, maltodextrin, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, povidone, silicon dioxide, sodium starch glycolate, stearic acid, talc, titanium dioxide

8335596

Dist. by: Reckitt Benckiser, Parsippany, NJ 07054-0224 ©2018 RB

060718



Drug Facts

Active ingredients (in each liquid gel)

Acetaminophen 325 mg.....Pain reliever/fever reducer
Dextromethorphan HBr 10 mg.....Cough suppressant
Phenylephrine HCl 5 mg.....Nasal decongestant

Purposes

Uses

- temporarily relieves these common cold and flu symptoms:
 - nasal congestion
 - cough
 - headache
 - minor aches and pains
 - sinus congestion and pressure
- temporarily reduces fever
- temporarily promotes nasal and/or sinus drainage

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 12 liquid gels in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks daily while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening ■ blisters ■ rash

If a skin reaction occurs, stop use and seek medical help right away.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's Disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- liver disease
- diabetes
- thyroid disease
- trouble urinating due to an enlarged prostate gland
- persistent or chronic cough such as occurs with smoking, asthma, or emphysema
- cough that occurs with too much phlegm (mucus) ▶
- heart disease
- high blood pressure

Drug Facts (continued)

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin

When using this product do not use more than directed

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back, or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- do not take more than directed (see Overdose warning)
- do not take more than 12 liquid gels in any 24-hour period
- adults and children 12 years of age and over: take 2 liquid gels every 4 hours
- children under 12 years of age: do not use

Other information

- store at 20-25°C (68-77°F)
- avoid excessive heat

Inactive ingredients

FD&C red no. 40, FD&C yellow no. 6, gelatin, glycerin, hypromellose, isopropyl alcohol, lecithin, light mineral oil, polyethylene glycol, povidone, propylene glycol, sorbitol sorbitan solution, titanium dioxide, water

Questions? 1-866-MUCINEX (1-866-682-4639)

You may also report side effects to this phone number.



CONGESTION &
HEADACHE

MAXIMUM STRENGTH

Maximum Strength per 4-hour dose
Do not take more than a total of
12 liquid gels in a 24-hour period.
Take only as directed.
Keep carton for full information.

PARENTS:
Learn about teen medicine abuse
www.StopMedicineAbuse.org



<p>Drug Facts</p> <p>Active ingredients Purposes (in each caplet)</p> <p>Mucinex FAST-MAX DAY TIME COLD & FLU</p> <p>Acetaminophen 325 mg.....Pain reliever/fever reducer</p> <p>Dextromethorphan HBr 10 mg.....Cough suppressant</p> <p>Guaifenesin 200 mg.....Expectorant</p> <p>Phenylephrine HCl 5 mg.....Nasal decongestant</p> <p>Active ingredients Purposes (in each caplet)</p> <p>Mucinex FAST-MAX</p>	<p>Drug Facts (continued)</p> <ul style="list-style-type: none"> ■ helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive (DAY TIME only) ■ controls cough to help you get to sleep ■ temporarily reduces fever <p>Warnings</p> <p>Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:</p> <ul style="list-style-type: none"> ■ more than 12 caplets in 24 hours, which is the maximum daily 	<p>Drug Facts (continued)</p> <p>Do not use</p> <ul style="list-style-type: none"> ■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. ■ with any other product containing diphenhydramine, even one used on the skin (NIGHT TIME only) ■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or medical conditions) 	<p>Drug Facts (continued)</p> <ul style="list-style-type: none"> ■ persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema ■ cough that occurs with too much phlegm (mucus) <p>Ask a doctor or pharmacist before use if you are</p> <ul style="list-style-type: none"> ■ taking the blood thinning drug warfarin ■ taking sedatives or tranquilizers (NIGHT TIME only) <p>When using this product</p> <ul style="list-style-type: none"> ■ do not use more than directed ■ excitability may occur
---	---	---	---

NIGHT TIME COLD & FLU

Acetaminophen 325 mg
Pain reliever/fever reducer
 Diphenhydramine HCl 12.5 mg
Antihistamine/cough suppressant
 Phenylephrine HCl 5 mgNasal decongestant

Uses

- temporarily relieves these common cold and flu symptoms:
 - cough
 - minor aches and pains
 - headache
 - nasal congestion
 - sore throat
 - sinus congestion and pressure
 - runny nose (**NIGHT TIME only**)
 - sneezing (**NIGHT TIME only**)
 - itching of the nose or throat (**NIGHT TIME only**)
 - itchy, watery eyes due to hay fever (**NIGHT TIME only**)

which is the maximum daily amount

- with other drugs containing acetaminophen
- 3 or more alcoholic drinks daily while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- liver disease
- heart disease
- diabetes
- high blood pressure
- thyroid disease
- trouble urinating due to an enlarged prostate gland
- glaucoma (**NIGHT TIME only**)
- a breathing problem such as emphysema or chronic bronchitis (**NIGHT TIME only**)

especially in children (**NIGHT TIME only**)

- marked drowsiness may occur (**NIGHT TIME only**)
- alcohol, sedatives, and tranquilizers may increase drowsiness (**NIGHT TIME only**)
- avoid alcoholic drinks (**NIGHT TIME only**)
- be careful when driving a motor vehicle or operating machinery (**NIGHT TIME only**)

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion, or cough gets worse or lasts more than 7 days

Drug Facts (continued)

- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back, or occurs with fever, rash, or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- do not take more than directed (see **Overdose warning**)
- do not take more than 12 caplets in any 24-hour period
- adults and children 12 years of age and over: take 2 caplets every 4 hours
- children under 12 years of age: do not use

Drug Facts (continued)**Other information**

- store between 20-25°C (68-77°F)

Inactive ingredients
(Mucinex FAST-MAX DAY TIME COLD & FLU)

corn starch, croscopovidone, FD&C red no. 40 aluminum lake, FD&C yellow no. 6 aluminum lake, magnesium stearate, maltodextrin, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, povidone, silicon dioxide, sodium starch glycolate, stearic acid, talc, titanium dioxide

Inactive ingredients
(Mucinex FAST-MAX NIGHT TIME COLD & FLU)

corn starch, croscarmellose sodium, croscopovidone, FD&C blue no. 1 aluminum lake, FD&C blue no. 2 aluminum lake, ferric oxide yellow, magnesium stearate, methacrylic acid-ethyl acrylate copolymer (1:1) type A, mica, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, povidone, silicon dioxide, sodium bicarbonate, stearic acid, talc, titanium dioxide

PEEL CORNER TO READ COMPLETE DRUG FACTS AND INFORMATION

Do not take the Mucinex Fast-Max DAY TIME COLD & FLU and Mucinex Fast-Max NIGHT TIME COLD & FLU caplets at the same time. Always wait at least 4 hours before taking another dose of Mucinex caplets. Do not take more than a total of 12 caplets in any 24-hour period.

Take only as directed.

102318 8351779



MAXIMUM STRENGTH

NDC 72854-138-66

Mucinex[®]

FAST[®] MAX[®]

KICKSTART[®]
SEVERE COLD & FLU

Acetaminophen » Pain Reliever/Fever Reducer

Dextromethorphan HBr » Cough Suppressant

Guaifenesin » Expectorant

Phenylephrine HCl » Nasal Decongestant

**COMEBACK
WITH A
KICK**



MENTHOL FLAVORED LIQUID

6 FL OZ (180 mL)

FOR AGES 12+

3259831

030823

Drug Facts

Active ingredients (in each 20 mL)

Purposes

Acetaminophen 650 mg	Pain reliever/fever reducer
Dextromethorphan HBr 20 mg	Cough suppressant
Guaifenesin 400 mg	Expectorant
Phenylephrine HCl 10 mg	Nasal decongestant

Uses

- temporarily relieves these common cold and flu symptoms:
 - cough
 - nasal congestion
 - minor aches and pains
 - sore throat
 - headache
 - stuffy nose
 - sinus congestion and pressure
- temporarily reduces fever
- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 6 doses in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks daily while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Drug Facts (continued)**Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- liver disease
- heart disease
- diabetes
- high blood pressure
- thyroid disease
- trouble urinating due to an enlarged prostate gland
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- cough that occurs with too much phlegm (mucus)

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin

When using this product do not use more than directed

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur

▶
PEEL HERE

Drug Facts (continued)

- cough comes back, or occurs with fever, rash, or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.


Keep out of reach of children.

Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- **do not take more than directed** (see **Overdose warning**)
- do not take more than 6 doses in any 24-hour period
- measure only with dosing cup provided
- do not use dosing cup with other products
- dose as follows or as directed by a doctor
- adults and children 12 years of age and over: 20 mL in dosing cup provided every 4 hours
- children under 12 years of age: do not use

Other information

- each 20 mL contains: **sodium 12 mg**
 - store at 20-25°C (68-77°F)
 - do not refrigerate
- 

**PEEL CORNER TO READ COMPLETE
DRUG FACTS AND INFORMATION**

Maximum Strength per 4-hour dose
Tamper evident: Do not use if neckband
on bottle cap is broken or missing.

*Helps to relieve these symptoms day or night

PARENTS:

Learn about teen medicine abuse

www.StopMedicineAbuse.org



3 63824 99696 7

Dist. by: RB Health (US)
Parsippany, NJ 07054-0224
©2023 RB Health
022423

LOT:

3259830

EXP:

MADE IN:



RB Artwork and Print Specification

Product Reference No.	8526090
2011 Ref	TR1518732
Brand	B
Category	Adult
Segment Group	Fast Max
Segment	All in One Night Time Cold & Flu
Print Size	400mm
Market Country	USA
Date	11/03/19

RB1 Contact: Anthony D'Amico

Artwork Type: IDM Commercial

Component Code (ID)	8377044
Parent Technical Packaging	8377044
Classification	8377044
Product Code (ID)	43024-04320-00
Country Code	USA & Canada
Language	EN
Language (Printed)	EN
Language (Packaging)	EN

Print Size: 400mm

Print Size: 400mm

Print Size: 400mm

Print Size: 400mm

Print Size: 400mm

Print Size: 400mm

Print Size: 400mm

Print Size: 400mm

Print Size: 400mm

Print Size: 400mm

Print Size: 400mm

Print Size: 400mm

Print Size: 400mm

Print Size: 400mm

Print Size: 400mm

Print Size: 400mm

Print Size: 400mm

Print Size: 400mm

Print Size: 400mm

Print Size: 400mm

Print Size: 400mm

Print Size: 400mm

Print Size: 400mm

Print Size: 400mm

Print Size: 400mm

Print Size: 400mm

Print Size: 400mm

Print Size: 400mm

Print Size: 400mm

Print Size: 400mm

Print Size: 400mm

Print Size: 400mm

Print Size: 400mm

Print Size: 400mm

Print Size: 400mm

Print Size: 400mm

Print Size: 400mm

Print Size: 400mm

Print Size: 400mm

Print Size: 400mm

Print Size: 400mm

Print Size: 400mm

Print Size: 400mm

Print Size: 400mm

Print Size: 400mm

Print Size: 400mm

Print Size: 400mm

Print Size: 400mm

Print Size: 400mm

Print Size: 400mm

Print Size: 400mm

Print Size: 400mm

Print Size: 400mm

Print Size: 400mm

Print Size: 400mm

Print Size: 400mm

Print Size: 400mm

Print Size: 400mm

Print Size: 400mm

Print Size: 400mm

Print Size: 400mm

Print Size: 400mm

Print Size: 400mm

Print Size: 400mm

Print Size: 400mm

Print Size: 400mm

Print Size: 400mm

Print Size: 400mm

Print Size: 400mm

Print Size: 400mm

Print Size: 400mm

BARCODE INFO

Barcode Type	UPC-A
Barcode Format	3-00000-000
Barcode Length	12
Barcode Height	23.2mm
Barcode Width	15.0mm
Barcode Depth	2.0mm
Barcode Color	Black

Barcode Color: Black

Barcode Color: Black

Barcode Color: Black

Barcode Color: Black

Barcode Color: Black

Barcode Color: Black

Barcode Color: Black

Barcode Color: Black

Barcode Color: Black

Barcode Color: Black

Barcode Color: Black

Barcode Color: Black

Barcode Color: Black

Barcode Color: Black

Barcode Color: Black

Barcode Color: Black

Barcode Color: Black

Barcode Color: Black

Barcode Color: Black

Barcode Color: Black

Barcode Color: Black

Barcode Color: Black

Barcode Color: Black

Barcode Color: Black

Barcode Color: Black

Barcode Color: Black

Barcode Color: Black

Barcode Color: Black

Barcode Color: Black

Barcode Color: Black

Barcode Color: Black

Barcode Color: Black

Barcode Color: Black

Barcode Color: Black

Barcode Color: Black

Barcode Color: Black

Barcode Color: Black

Barcode Color: Black

Barcode Color: Black

Barcode Color: Black

Barcode Color: Black

Barcode Color: Black

Barcode Color: Black

Barcode Color: Black

Barcode Color: Black

Barcode Color: Black

Barcode Color: Black

Barcode Color: Black

Barcode Color: Black

Barcode Color: Black

Barcode Color: Black

Barcode Color: Black

Barcode Color: Black

MIN PT SIZE & HEIGHT (Non Automated)	
Text Size: 12pt	12pt
Text Size: 10pt	10pt
Text Size: 8pt	8pt
Text Size: 6pt	6pt
Text Size: 4pt	4pt
Text Size: 3pt	3pt
Text Size: 2pt	2pt
Text Size: 1pt	1pt
Text Size: 0.5pt	0.5pt
Text Size: 0.25pt	0.25pt
Text Size: 0.125pt	0.125pt
Text Size: 0.0625pt	0.0625pt
Text Size: 0.03125pt	0.03125pt
Text Size: 0.015625pt	0.015625pt
Text Size: 0.0078125pt	0.0078125pt
Text Size: 0.00390625pt	0.00390625pt
Text Size: 0.001953125pt	0.001953125pt
Text Size: 0.0009765625pt	0.0009765625pt
Text Size: 0.00048828125pt	0.00048828125pt
Text Size: 0.000244140625pt	0.000244140625pt
Text Size: 0.0001220703125pt	0.0001220703125pt
Text Size: 0.00006103515625pt	0.00006103515625pt
Text Size: 0.000030517578125pt	0.000030517578125pt
Text Size: 0.0000152587890625pt	0.0000152587890625pt
Text Size: 0.00000762939453125pt	0.00000762939453125pt
Text Size: 0.000003814697265625pt	0.000003814697265625pt
Text Size: 0.0000019073486328125pt	0.0000019073486328125pt
Text Size: 0.00000095367431640625pt	0.00000095367431640625pt
Text Size: 0.000000476837158203125pt	0.000000476837158203125pt
Text Size: 0.0000002384185791015625pt	0.0000002384185791015625pt
Text Size: 0.00000011920928955078125pt	0.00000011920928955078125pt
Text Size: 0.000000059604644775390625pt	0.000000059604644775390625pt
Text Size: 0.0000000298023223876953125pt	0.0000000298023223876953125pt
Text Size: 0.00000001490116119384765625pt	0.00000001490116119384765625pt
Text Size: 0.000000007450580596923828125pt	0.000000007450580596923828125pt
Text Size: 0.0000000037252902984619140625pt	0.0000000037252902984619140625pt
Text Size: 0.00000000186264514923095703125pt	0.00000000186264514923095703125pt
Text Size: 0.000000000931322574615478515625pt	0.000000000931322574615478515625pt
Text Size: 0.0000000004656612873077392578125pt	0.0000000004656612873077392578125pt
Text Size: 0.00000000023283064365386962890625pt	0.00000000023283064365386962890625pt
Text Size: 0.000000000116415321826934814453125pt	0.000000000116415321826934814453125pt
Text Size: 0.0000000000582076609134674072265625pt	0.0000000000582076609134674072265625pt
Text Size: 0.00000000002910383045673370361328125pt	0.00000000002910383045673370361328125pt
Text Size: 0.000000000014551915228366851806640625pt	0.000000000014551915228366851806640625pt
Text Size: 0.0000000000072759576141834259033203125pt	0.0000000000072759576141834259033203125pt
Text Size: 0.00000000000363797880709171295166015625pt	0.00000000000363797880709171295166015625pt
Text Size: 0.000000000001818989403545856475830078125pt	0.000000000001818989403545856475830078125pt
Text Size: 0.0000000000009094947017729282379150390625pt	0.0000000000009094947017729282379150390625pt
Text Size: 0.00000000000045474735088641211895751953125pt	0.00000000000045474735088641211895751953125pt
Text Size: 0.000000000000227373675443206059478759765625pt	0.000000000000227373675443206059478759765625pt
Text Size: 0.0000000000001136868377216030297393798828125pt	0.0000000000001136868377216030297393798828125pt
Text Size: 0.00000000000005684341886080151486968994140625pt	0.00000000000005684341886080151486968994140625pt
Text Size: 0.000000000000028421709430400757434844970703125pt	0.000000000000028421709430400757434844970703125pt
Text Size: 0.0000000000000142108547152003787174224853515625pt	0.0000000000000142108547152003787174224853515625pt
Text Size: 0.000000000000007105427357600189358711224267578125pt	0.000000000000007105427357600189358711224267578125pt
Text Size: 0.0000000000000035527136788000946793556121337890625pt	0.0000000000000035527136788000946793556121337890625pt
Text Size: 0.0000000000000017763568394000473396778060668953125pt	0.0000000000000017763568394000473396778060668953125pt
Text Size: 0.00000000000000088817841970002366983890303344765625pt	0.00000000000000088817841970002366983890303344765625pt
Text Size: 0.000000000000000444089209850011834919451516723828125pt	0.000000000000000444089209850011834919451516723828125pt
Text Size: 0.0000000000000002220446049250059174597257583619140625pt	0.0000000000000002220446049250059174597257583619140625pt
Text Size: 0.00000000000000011102230246250295872986287918095703125pt	0.00000000000000011102230246250295872986287918095703125pt
Text Size: 0.0000000000000000555111512312514793649314395904765625pt	0.0000000000000000555111512312514793649314395904765625pt
Text Size: 0.00000000000000002775557561562573968246571979523828125pt	0.00000000000000002775557561562573968246571979523828125pt
Text Size: 0.000000000000000013877787807812869841232859897619140625pt	0.000000000000000013877787807812869841232859897619140625pt
Text Size: 0.0000000000000000069388939039064349206164299488095703125pt	0.0000000000000000069388939039064349206164299488095703125pt
Text Size: 0.000000000000000003469446951953217460308214974404765625pt	0.000000000000000003469446951953217460308214974404765625pt
Text Size: 0.0000000000000000017347234759766087301541074872023828125pt	0.0000000000000000017347234759766087301541074872023828125pt
Text Size: 0.00000000000000000086736173798830436507705374360119140625pt	0.00000000000000000086736173798830436507705374360119140625pt
Text Size: 0.000000000000000000433680868994152182538526871800595703125pt	0.000000000000000000433680868994152182538526871800595703125pt
Text Size: 0.00000000000000000021684043449707609126926343590029765625pt	0.00000000000000000021684043449707609126926343590029765625pt
Text Size: 0.000000000000000000108420217248538045634631717950148828125pt	0.000000000000000000108420217248538045634631717950148828125pt
Text Size: 0.0000000000000000000542101086242690228173158589750744140625pt	0.0000000000000000000542101086242690228173158589750744140625pt
Text Size: 0.00000000000000000002710505431213451140865792948753720703125pt	0.00000000000000000002710505431213451140865792948753720703125pt
Text Size: 0.000000000000000000013552527156067255704328964743768603515625pt	0.000000000000000000013552527156067255704328964743768603515625pt
Text Size: 0.0000000000000000000067762635780336278521644823721843017578125pt	0.0000000000000000000067762635780336278521644823721843017578125pt
Text Size: 0.00000000000000000000338813178901681392608224118609215087890625pt	0.00000000000000000000338813178901681392608224118609215087890625pt
Text Size: 0.000000000000000000001694065894508406963041120593046075439453125pt	0.000000000000000000001694065894508406963041120593046075439453125pt
Text Size: 0.0000000000000000000008470329472542034815205602965230377197265625pt	0.0000000000000000000008470329472542034815205602965230377197265625pt
Text Size: 0.00000000000000000000042351647362710174076028014826151885986328125pt	0.00000000000000000000042351647362710174076028014826151885986328125pt
Text Size: 0.000000000000000000000211758236813550870380140074130759429931640625pt	0.000000000000000000000211758236813550870380140074130759429931640625pt
Text Size: 0.0000000000000000000001058791184067754351900700370653797149658203125pt	0.0000000000000000000001058791184067754351900700370653797149658203125pt
Text Size: 0.00000000000000000000005293955920338771759503501853268985748291015625pt	0.00000000000000000000005293955920338771759503501853268985748291015625pt
Text Size: 0.000000000000000000000026469779601693858797517509266344928741455078125pt	0.000000000000000000000026469779601693858797517509266344928741455078125pt
Text Size: 0.0000000000000000000000132348898008469293987587546331724643707275390625pt	0.0000000000000000000000132348898008469293987587546331724643707275390625pt
Text Size: 0.00000000000000000000000661744490042346469937937731658623218536376953125pt	0.00000000000000000000000661744490042346469937937731658623218536376953125pt
Text Size: 0.000000000000000000000003308722450211732349689688658293116092681884765625pt	0.000000000000000000000003308722450211732349689688658293116092681884765625pt
Text Size: 0.0000000000000000000000016543612251058661748448443291465580463409423828125pt	0.0000000000000000000000016543612251058661748448443291465580463409423828125pt
Text Size: 0.00000000000000000000000082718061255293308742242216457327902317047119140625pt	0.00000000000000000000000082718061255293308742242216457327902317047119140625pt
Text Size: 0.000000000000000000000000413590306276466543711211082286639511585235595703125pt	0.000000000000000000000000413590306276466543711211082286639511585235595703125pt
Text Size: 0.00000000000000000000000020679515313823327185560554114331975792761779787500	

MAXIMUM STRENGTH



Tamper evident: Do not use if carton is damaged or if printed seal on blister is broken or missing.

**SEVERE CONGESTION
& COUGH**

NDC 63824-193-21

MAXIMUM STRENGTH



**SEVERE CONGESTION
& COUGH**

Dextromethorphan HBr – Cough Suppressant
Guaifenesin – Expectorant
Phenylephrine HCl – Nasal Decongestant

- ✓ Controls Cough
- ✓ Relieves Nasal & Chest Congestion
- ✓ Thins & Loosens Mucus

Actual Size

20 CAPLETS

FOR AGES 12+



HEALTH • HYGIENE • HOME

www.mucinex.com
Patents: www.rb.com/patents

Dist. by: RB Health (US)
Parsippany, NJ 07054-0224
©2020 RB Health

041320 3141649



3 63824 19320 5

LOT:

EXP:

MADE IN:
3141649



Drug Facts

Active ingredients (in each caplet)

Dextromethorphan HBr 10 mg..... Cough suppressant
Guaifenesin 200 mg..... Expectorant
Phenylephrine HCl 5 mg..... Nasal decongestant

Purposes

Uses

- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive
- temporarily relieves:
 - cough due to minor throat and bronchial irritation as may occur with the common cold or inhaled irritants
 - the intensity of coughing
 - the impulse to cough to help you get to sleep
 - nasal congestion due to a cold

Warnings

Do not use

- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- heart disease
- diabetes
- high blood pressure
- thyroid disease
- trouble urinating due to an enlarged prostate gland
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- cough that occurs with too much phlegm (mucus) ▶

Drug Facts (continued)

When using this product do not use more than directed

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- symptoms do not get better within 7 days or occur with fever
- cough comes back, or occurs with fever, rash, or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- do not take more than 12 caplets in any 24-hour period
- adults and children 12 years of age and over: take 2 caplets every 4 hours
- children under 12 years of age: do not use

Other information

- store at 20-25°C (68-77°F)

Inactive ingredients

croscarmellose sodium, FD&C blue no. 2 aluminum lake, FD&C red no. 40 aluminum lake, methacrylic acid-ethyl acrylate copolymer, mica, microcrystalline cellulose, polyethylene glycol 3350, polysorbate 80, polyvinyl alcohol, povidone K29/32, sodium bicarbonate, talc, titanium dioxide

Questions? 1-866-MUCINEX (1-866-682-4639)

You may also report side effects to this phone number.



MAXIMUM STRENGTH

**SEVERE CONGESTION
& COUGH**

Maximum Strength per 4-hour dose
Do not take more than a total of 12 caplets
in any 24-hour period.
Take only as directed.
Keep carton for full information.



8-1715-697-09
ORG

8-1715-697-09
ORG

Drug Facts

Active ingredients (in each caplet)

Dextromethorphan HBr 10 mg.....Cough suppressant
Guaifenesin 200 mg.....Expectorant
Phenylephrine HCl 5 mg.....Nasal decongestant

Purposes

Uses

- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive
- temporarily relieves:
 - cough due to minor throat and bronchial irritation as may occur with the common cold or inhaled irritants
 - the intensity of coughing
 - the impulse to cough to help you get to sleep
 - nasal congestion due to a cold

Warnings

Do not use

- If you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- heart disease
- diabetes
- high blood pressure
- thyroid disease
- trouble urinating due to an enlarged prostate gland
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- cough that occurs with too much phlegm (mucus) ▶

Drug Facts (continued)

When using this product do not use more than directed

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- symptoms do not get better within 7 days or occur with fever
- cough lasts more than 7 days, comes back, or occurs with fever, rash, or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- do not take more than 12 caplets in any 24-hour period
- adults and children 12 years of age and over: take 2 caplets every 4 hours
- children under 12 years of age: do not use

Other information

- store at 20-25°C (68-77°F)

Inactive ingredients colloidal silicon dioxide, FD&C blue no. 2 aluminum lake, FD&C red no. 40 aluminum lake, magnesium stearate, maltodextrin, mica, microcrystalline cellulose, polyethylene glycol, polysorbate 80, polyvinyl alcohol, povidone, pregelatinized starch, sodium starch glycolate, stearic acid, talc, titanium dioxide

Questions? 1-866-MUCINEX (1-866-682-4639)



MIN PT SIZE & HEIGHT (Non-Automated)			
NET WT (6.5 FL OZ 190ml):	.10inch	10pt	
Drug Facts Title:	.06inch	XXpt	
Active Ingredient Title:	.07inch	7pt	
Active Ingredient (BOP Ingredient):	.06inch	XXpt	
Min Size of all other Copy:	.060inch	6.6pt	
<div> <div>This Box 1/2 x 1/2 inch Square</div> <div> Front PDP in square inches : 12.136 Rear PDP in square inches : XX.XX </div> </div>			

RB Artwork and Print Specification																	
Trident Reference No:	RB962073																
ZEN Ref:	TR1237086																
Brand:	A																
Category:	Mucinex																
Segment Group:	Adult																
Segment:	Fast Max																
Pack Size:	Severe Congestion Et Cough																
Market/Country:	611 oz																
Date:	USA																
RBH Contact:	17/03/17																
RBH Contact: Laura Feld																	
Artwork Type: IDM Commercial																	
Component Code (if applicable):	3048243																
Parent Technical Packaging Specification:	D0319825																
Finished Goods Code:	63624-01485-10																
Supply Point:	RB Hull																
Pharmacoce No/NE:	N/A																
Edgemark Position:	NA																
CAD Cam Ref:	Muc-Lbl-D0319825-90x87mm																
Printer:	Multi Packaging Solutions (Bourne, UK)																
Substrate:	Paper White																
Technical Et Non Printing Items																	
<input type="checkbox"/> Cutter <input type="checkbox"/> Outlets	<input type="checkbox"/> Cutter 2 (if applicable) <input type="checkbox"/> Outlets 2 (if applicable)																
Colours (Label Front)																	
<table border="1"> <tr> <td>PRINTING SWK 1</td><td>PRINTING SWK 2</td><td>PRINTING SWK 3</td><td>PRINTING SWK 4</td></tr> <tr> <td>1 FLX</td><td>2 FLX</td><td>3 FLX</td><td>4 FLX</td></tr> </table>	PRINTING SWK 1	PRINTING SWK 2	PRINTING SWK 3	PRINTING SWK 4	1 FLX	2 FLX	3 FLX	4 FLX	<table border="1"> <tr> <td>PRINTING SWK 5</td><td>PRINTING SWK 6</td><td>PRINTING SWK 7</td><td>PRINTING SWK 8</td></tr> <tr> <td>5 FLX</td><td>6 FLX</td><td>7 FLX</td><td>8 FLX</td></tr> </table>	PRINTING SWK 5	PRINTING SWK 6	PRINTING SWK 7	PRINTING SWK 8	5 FLX	6 FLX	7 FLX	8 FLX
PRINTING SWK 1	PRINTING SWK 2	PRINTING SWK 3	PRINTING SWK 4														
1 FLX	2 FLX	3 FLX	4 FLX														
PRINTING SWK 5	PRINTING SWK 6	PRINTING SWK 7	PRINTING SWK 8														
5 FLX	6 FLX	7 FLX	8 FLX														
Barcode Info																	
Barcode Type:	N/A																
Barcode Number:	N/A																
Magnification:	N/A																
Truncated By:	N/A																
Full Height:	N/A																
Bar Height (Smallest Bar):	N/A																
BWR:	N/A																
Encoded Data:	N/A																
 HEALTH • HYGIENE • HOME																	
 TRIDENT																	
Connaught House, Connaught Road, Kingswood Business Park, Hull, HU7 3AP, England. T: +44 (0) 1482 828100																	
Please note that any low resolution paper Canon colour copies associated with this job should be referred to for content, layout and colour separation only.																	
STUDIO USE ONLY	Adam Simmons v2.0																
SimArt check results: G-1; Q-0; R-0; - AS - 03/17/17 15:26:48																	
ARTWORK APPROVAL																	
Approved by Sonoco Trident on behalf of Recidtt Benckiser																	
Signed: _____																	
Date: _____																	
NOT APPROVED FOR COLOUR																	
UNDER NO CIRCUMSTANCES SHOULD THIS ARTWORK BE ALTERED WITHOUT PRIOR PERMISSION FROM TRIDENT.																	



Drug Facts

Active Ingredients (in each 20 mL)

Dextromethorphan HBr 20 mg
.....Cough suppressant
Guaifenesin 400 mg.....Expectorant
Phenylephrine HCl 10 mg
.....Nasal decongestant

Uses

- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive
- temporarily relieves:
 - cough due to minor throat and bronchial irritation as may occur with the common cold or inhaled irritants
 - the intensity of coughing
 - nasal congestion due to a cold
 - temporarily helps you cough less

Warnings

- Do not use**
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or

Drug Facts (continued)

emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- heart disease
- high blood pressure
- thyroid disease
- diabetes
- trouble urinating due to an enlarged prostate gland
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- cough that occurs with too much phlegm (mucus)

When using this product do not use more than directed

- Stop use and ask a doctor if**
- nervousness, dizziness, or sleeplessness occur
 - symptoms do not get better within 7 days or occur with fever
 - cough comes back, or occurs with rash or headache that

Drug Facts (continued)

lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- do not take more than 6 doses in any 24-hour period
- measure only with dosing cup provided
- do not use dosing cup with other products
- dose as follows or as directed by a doctor
- adults and children 12 years of age and older: 20 mL in dosing cup provided every 4 hours
- children under 12 years of age: do not use

Other information

- each 20 mL contains: sodium 9 mg
- store between 20-25°C (68-77°F)
- dosing cup provided
- do not refrigerate

PEEL CORNER TO READ COMPLETE DRUG FACTS AND INFORMATION

Tamper evident: Do not use if neckband on bottle cap is broken or missing.

PARENTS:

Learn about teen medicine abuse

www.StopMedicineAbuse.org



3 63824 54166 2

LOT:

EXP:

3027786

Drug Facts (continued)

Inactive ingredients

anhydrous citric acid, D&C yellow no.10, edetate disodium,

3005
Fruite, blue 3005, glycerin,
propyl gallate, propylene glycol,
sodium benzoate, sodium citrate,
sorbitol, sucralose, water,
xanthan gum

Questions?**1-866-MUCINEX (1-866-682-4639)**

You may also report side effects
to this phone number.

**HEALTH • HYGIENE • HOME**

Please visit our web site
www.mucinex.com

Dist. by: Reckitt Benckiser
Parsippany, NJ 07054-0224
Made in England ©2016 RB



**PEEL CORNER TO READ COMPLETE
DRUG FACTS AND INFORMATION**

Maximum Strength per 4-hour dose

Tamper evident: Do not use
if neckband on bottle cap
is broken or missing.

Drug Facts

**Active
ingredients
(in each 20 mL)**

Dextromethorphan HBr 20 mg

Cough suppressant

Purposes

PARENTS!
Learn about teen medicine abuse
www.StopMedicineAbuse.org



3 63824 54166 2

LOT: 3087260

EXP:

3007.....Cough suppressant
Guaifenesin 400 mg.....Expectorant
Phenylephrine HCl 10 mg
.....Nasal decongestant

Uses

- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive
- temporarily relieves:
 - cough due to minor throat and bronchial irritation as may occur with the common cold or inhaled irritants
 - the intensity of coughing
 - nasal congestion due to a cold
 - temporarily helps you cough less

Warnings

Do not use

- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or

Drug Facts (continued)

emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- heart disease
- high blood pressure
- thyroid disease
- diabetes
- trouble urinating due to an enlarged prostate gland
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- cough that occurs with too much phlegm (mucus)

When using this product do not use more than directed

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- symptoms do not get better within 7 days or occur

Drug Facts (continued)

lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- do not take more than 6 doses in any 24-hour period
- measure only with dosing cup provided
- do not use dosing cup with other products
- dose as follows or as directed by a doctor
- adults and children 12 years of age and older: 20 mL in dosing cup provided every 4 hours
- children under 12 years of age: do not use

Other information

- each 20 mL contains:
sodium 9 mg
- store at 20-25°C

with fever
■ cough comes back, or occurs
with rash or headache that ▶

(68-77°F)
■ dosing cup provided
■ do not refrigerate ▶

Drug Facts (continued)

Inactive ingredients

anhydrous citric acid, D&C
yellow no.10, edetate disodium,
FD&C blue no.1, flavors, glycerin,
propyl gallate, propylene glycol,
sodium benzoate, sodium citrate,
sorbitol, sucralose, water,
xanthan gum

Questions?

1-866-MUCINEX (1-866-682-4639)
You may also report side effects
to this phone number.



HEALTH ▶ HYGIENE ▶ HOME

Please visit our website
www.mucinex.com

Patents: www.rb.com/patents

Dist. by: RB Health (US)
Parsippany, NJ 07054-0224
Made in England ©2019 RB Health



Drug Facts

Active ingredients (in each 20 mL)

Purposes

Acetaminophen 650 mg	Pain reliever/fever reducer
Dextromethorphan HBr 20 mg	Cough suppressant
Phenylephrine HCl 10 mg	Nasal decongestant
Triprolidine HCl 2.5 mg	Antihistamine

Uses

- temporarily relieves these common cold and flu symptoms:
 - cough
 - minor aches and pains
 - headache
 - sinus congestion and pressure
 - sneezing
 - nasal congestion
 - sore throat
 - runny nose

Drug Facts (continued)

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a

- itching of the nose or throat
- itchy, watery eyes due to hay fever
- temporarily reduces fever
- controls cough to help you get to sleep

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 4000 mg in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks daily while using this product

your prescription drug contains an opioid, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- liver disease
- heart disease
- diabetes
- high blood pressure
- thyroid disease
- glaucoma
- trouble urinating due to an enlarged prostate gland
- a breathing problem such as emphysema or chronic bronchitis
- persistent or chronic cough such as occurs with smoking, asthma, or emphysema
- cough that occurs with too much phlegm (mucus)

PEEL HERE

Drug Facts (continued)

Ask a doctor or pharmacist before use if you are

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers

When using this product

- **do not use more than directed**
- excitability may occur, especially in children
- marked drowsiness may occur
- alcohol, sedatives, and tranquilizers may increase drowsiness
- avoid alcoholic drinks
- use caution when driving a motor vehicle or operating machinery

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back, or occurs with fever, rash, or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Drug Facts (continued)

Directions

- **do not take more than directed (see Overdose warning)**
- do not take more than 4 doses in any 24-hour period
- measure only with dosing cup provided
- do not use dosing cup with other products
- dose as follows or as directed by a doctor
- adults and children 12 years of age and over: 20 mL in dosing cup provided every 4 hours
- children under 12 years of age: do not use

Other information

- each 20 mL contains: sodium 12 mg
- store at 20-25°C (68-77°F)
- do not refrigerate

Inactive ingredients ammonium glycyrrhizate, anhydrous citric acid, cochineal carmine, edetate disodium, flavors, glycerin (soy), maltodextrin, propylene glycol, purified water, sodium benzoate, sorbitol, sucralose, xanthan gum

Questions? 1-866-MUCINEX (1-866-682-4639)

You may also report side effects to this phone number.



HEALTH • HYGIENE • HOME

Patents:
www.rb.com/patents
Please visit our website
www.mucinex.com

**PEEL CORNER TO READ COMPLETE
DRUG FACTS AND INFORMATION**

Maximum Strength per 4-hour dose
**Tamper evident: Do not use if neckband
on bottle cap is broken or missing.**

121219

**FREE
FROM**

DATA MATRIX
B-1715-719-09
ORG

Drug Facts

Active ingredients (in each caplet)

Acetaminophen 162.5 mg...Pain reliever/fever reducer
Dextromethorphan HBr 5 mg...Cough suppressant
Guafenesin 100 mg...Expectorant
Phenylephrine HCl 2.5 mg...Nasal decongestant

Purposes

Uses

- temporarily relieves these common cold and flu symptoms:
 - cough
 - nasal congestion
 - headache
 - sore throat
 - minor aches and pains
 - sinus congestion and pressure
- temporarily reduces fever
- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive

Warnings

Liver warnings: This product contains acetaminophen. Severe liver damage may occur if your child takes:

- more than 5 doses in 24 hours, which is the maximum daily amount.

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- hives
- rash

 If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

in a child who is taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your child's prescription drug contains an MAOI, ask a doctor or pharmacist before giving this product.

Ask a doctor before use if the child has

- liver disease
- heart disease
- high blood pressure
- thyroid disease

Drug Facts (continued)

- diabetes
- persistent or chronic cough such as occurs with asthma
- cough that occurs with too much phlegm (mucus)

Ask a doctor or pharmacist before use if the child is taking the blood thinning drug warfarin

When using this product do not use more than directed (see Overdose warning)

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion, or cough gets worse or lasts more than 5 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back, or occurs with fever, rash, or headache that lasts. These could be signs of a serious condition.

Keep out of reach of children.

Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical even if you do not notice any signs or symptoms.

Directions

- this product does not contain directions or complete warnings for adult use
- do not give more than directed (see Overdose warning)
- dose as follows or as directed by a doctor
- children 6 to under 12 years of age: give 2 caplets every 4 hours while symptoms last; do not give more than 5 doses in any 24-hour period
- children under 6 years of age: do not use

Other information

- Store at 20-25°C (68-77°F)

Inactive ingredients

corn starch, croscopolone, FD&C blue no. 1 aluminum lake, magnesium stearate, maltodextrin, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, povidone, silicon dioxide, sodium starch glycolate, stearic acid, talc, titanium dioxide

Questions? 1-866-MUCINEX (1-866-682-4639)

You may also report side effects to this phone number.

PARENTS:

Learn about teen medicine abuse
www.StopMedicineAbuse.org

is an OTC brand in the Children's Cough/Cold category (Alphametrics/ProVoice Survey) *as compared to Mucinex caplets

Helps to relieve these symptoms day or night if your child has difficulty swallowing solid dosage forms, try our Mucinex Children's liquids or mini-melts products. Keep carton for full information.

B-1715-719-09
ORG

Mucinex
Junior
COLD & FLU

Tamper evident: Do not use if carton is damaged or if printed seal on blister is broken or missing.

Pediatrician Recommended

NUC 03824-070-20

NEW

Mucinex
Junior
COLD & FLU

Acetaminophen 162.5 mg - Pain Reliever/Fever Reducer
Dextromethorphan HBr 5 mg - Cough Suppressant
Guafenesin 100 mg - Expectorant
Phenylephrine HCl 2.5 mg - Nasal Decongestant

Ages 6+ yrs



Actual Size

20 CAPLETS

DO NOT CHEW

DATE
LOT & EXP.



www.mucinex.com

Dist. by: RB Health (US)
Paramus, NJ 07654-0224
©2019 RB Health
031819 3098705

HEALTH • HYGIENE • HOME
US Patents: www.rb.com/patents



Drug Facts

Active ingredients (in each 20 mL)

Purposes

Acetaminophen 650 mg... Pain reliever/fever reducer
Dextromethorphan HBr 20 mg... Cough suppressant
Phenylephrine HCl 10 mg... Nasal decongestant
Tripolidine HCl 2.5 mg... Antihistamine

Uses

- temporarily relieves these common cold and flu symptoms:
 - cough
 - nasal congestion
 - minor aches and pains
 - sore throat
 - headache
 - sinus congestion and pressure
 - runny nose
 - sneezing
 - itching of the nose or throat
 - itchy, watery eyes due to hay fever
- temporarily reduces fever
- controls cough to help you get to sleep

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 4,000 mg in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks daily while using this product

Drug Facts (continued)

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
 - blisters
 - rash
- If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- liver disease
- heart disease
- diabetes
- high blood pressure
- thyroid disease
- glaucoma
- trouble urinating due to an enlarged prostate gland
- a breathing problem such as emphysema or chronic bronchitis
- persistent or chronic cough such as occurs with smoking, asthma, or emphysema
- cough that occurs with too much phlegm (mucus)

PEEL HERE

Drug Facts (continued)

Ask a doctor or pharmacist before use if you are

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers

When using this product

- **do not use more than directed**
- excitability may occur, especially in children
- marked drowsiness may occur
- alcohol, sedatives, and tranquilizers may increase drowsiness
- avoid alcoholic drinks
- use caution when driving a motor vehicle or operating machinery

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back, or occurs with fever, rash, or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Drug Facts (continued)

Directions

- **do not take more than directed** (see **Overdose warning**)
- do not take more than 4 doses in any 24-hour period
- measure only with dosing cup provided
- do not use dosing cup with other products
- dose as follows or as directed by a doctor
- adults and children 12 years of age and over: 20 mL in dosing cup provided every 4 hours
- children under 12 years of age: do not use

Other information

- each 20 mL contains: **sodium 16 mg**
- store at 20-25°C (68-77°F)
- do not refrigerate

Inactive ingredients ammonium glycyrrhizate, anhydrous citric acid, ascorbic acid, edetate disodium, FD&C blue no. 1, FD&C red no. 40, flavors, glycerin (soy), propylene glycol, sodium benzoate, sorbitol, sucralose, triacetin, triethyl citrate, water, xanthan gum

Questions?

1-866-MUCINEX (1-866-682-4639)

Scan for FAQs and instructions on proper disposal of medicines



Please visit our website
www.mucinex.com

Patents:
www.reckitt.com/patents

NDC 63824-501-66



Mucinex[®]

NIGHTSHIFT

**COLD & FLU
CLEAR & COOL**

Acetaminophen – Pain Reliever/Fever Reducer
Dextromethorphan HBr – Cough Suppressant
Phenylephrine HCl – Nasal Decongestant
Triprolidine HCl – Antihistamine



NIGHT TIME

RELIEF FOR A BETTER

MORNING

✓ COUGH ✓ FEVER ✓ SORE THROAT
 ✓ RUNNY NOSE ✓ SNEEZING ✓ NASAL CONGESTION

3094779
6 FL OZ (180 mL)
FOR AGES 12+
030619

Drug Facts

Active ingredients (in each 20 mL)	Purposes
Acetaminophen 650 mg.....Pain reliever/fever reducer	
Dextromethorphan HBr 20 mg.....Cough suppressant	
Phenylephrine HCl 10 mg.....Nasal decongestant	
Triprolidine HCl 2.5 mg.....Antihistamine	

Uses

- temporarily relieves these common cold and flu symptoms:
 - cough
 - minor aches and pains
 - headache
 - sneezing
 - sinus congestion and pressure
 - itching of the nose or throat
 - itchy, watery eyes due to hay fever
- temporarily reduces fever
- controls cough to help you get to sleep

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 4000 mg in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks daily while using this product

Drug Facts (continued)

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- liver disease
- diabetes
- thyroid disease
- trouble urinating due to an enlarged prostate gland
- a breathing problem such as emphysema or chronic bronchitis
- persistent or chronic cough such as occurs with smoking, asthma, or emphysema
- cough that occurs with too much phlegm (mucus)
- heart disease
- high blood pressure
- glaucoma

PEEL HERE

Drug Facts (continued)

Ask a doctor or pharmacist before use if you are

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers

When using this product

- do not use more than directed
- excitability may occur, especially in children
- marked drowsiness may occur
- alcohol, sedatives, and tranquilizers may increase drowsiness
- avoid alcoholic drinks
- use caution when driving a motor vehicle or operating machinery

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- pain, nasal decongestion, or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back, or occurs with fever, rash, or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Drug Facts (continued)

Directions

- do not take more than directed (see Overdose warning)
- do not take more than 4 doses in any 24-hour period
- measure only with dosing cup provided
- do not use dosing cup with other products
- dose as follows or as directed by a doctor
- adults and children 12 years of age and over: 20 mL in doses as directed every 4 hours

**PEEL CORNER TO READ COMPLETE
DRUG FACTS AND INFORMATION**

Tamper evident: Do not use if neckband on bottle cap is broken or missing.

Maximum Strength per 4-hour dose

PARENTS:

Learn about teen medicine abuse

www.StopMedicineAbuse.org




NEW

**POWERFUL
NIGHTTIME
RELIEF**

031419







Drug Facts

Active ingredients (in each 20 mL)	Purposes
Acetaminophen 650 mg...Pain reliever/fever reducer	
Dextromethorphan HBr 20 mg.....Cough suppressant	
Phenylephrine HCl 10 mg.....Nasal decongestant	
Triprolidine HCl 2.5 mg.....Antihistamine	

Uses

- temporarily relieves these common cold and flu symptoms:
 - cough
 - minor aches and pains
 - headache
 - sinus congestion and pressure
 - runny nose
 - sneezing
 - itching of the nose or throat
 - itchy, watery eyes due to hay fever
- temporarily reduces fever
- controls cough to help you get to sleep

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 4000 mg in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks daily while using this product

Drug Facts (continued)

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- liver disease
- diabetes
- thyroid disease
- trouble urinating due to an enlarged prostate gland
- a breathing problem such as emphysema or chronic bronchitis
- persistent or chronic cough such as occurs with smoking, asthma, or emphysema
- cough that occurs with too much phlegm (mucus)
- heart disease
- high blood pressure
- glaucoma

PEEL HERE

Drug Facts (continued)

Ask a doctor or pharmacist before use if you are

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers

When using this product

- do not use more than directed
- excitability may occur, especially in children
- marked drowsiness may occur
- alcohol, sedatives, and tranquilizers may increase drowsiness
- avoid alcoholic drinks
- use caution when driving a motor vehicle or operating machinery

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back, or occurs with fever, rash, or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Drug Facts (continued)

Directions

- do not take more than directed (see Overdose warning)
- do not take more than 4 doses in any 24-hour period
- measure only with dosing cup provided
- do not use dosing cup with other products
- dose as follows or as directed by a doctor
- adults and children 12 years of age and over: 20 mL in dosing cup provided every 4 hours
- children under 12 years of age: do not use

Other information


- each 20 mL contains: sodium 16 mg
- store at 20-25°C (68-77°F)
- do not refrigerate

Inactive ingredients ammonium glycyrrhizate, anhydrous citric acid, ascorbic acid, edetate disodium, FD&C blue no. 1, FD&C red no. 40, flavors, glycerin (soy), propylene glycol, sodium benzoate, sorbitol, sucralose, triacetin, triethyl citrate, water, xanthan gum

Questions?

1.866.MUSICINEY (1.866.682.4630)

3019





3 63824 01473 9

Please visit our website
www.mucinex.com

Dist. by: RB Health (US)
Parsippany, NJ 07054-0224
©2019 RB Health
031119

LOT: 3094781

EXP:

MADE IN:

- temporarily relieves these common cold and flu symptoms:
- cough
- minor aches and pains
- headache
- sneezing
- sinus congestion and pressure
- itching of the nose or throat
- itchy, watery eyes due to hay fever
- temporarily reduces fever
- controls cough to help you get to sleep
- nasal congestion
- sore throat
- runny nose

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 4000 mg in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks daily while using this product

Drug Facts (continued)

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- liver disease
- heart disease
- diabetes
- high blood pressure
- thyroid disease
- glaucoma
- trouble urinating due to an enlarged prostate gland
- a breathing problem such as emphysema or chronic bronchitis
- persistent or chronic cough such as occurs with smoking, asthma, or emphysema
- cough that occurs with too much phlegm (mucus)

PEEL HERE

Drug Facts (continued)

Ask a doctor or pharmacist before use if you are

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers

When using this product

- do not use more than directed
- excitability may occur, especially in children
- marked drowsiness may occur
- alcohol, sedatives, and tranquilizers may increase drowsiness
- avoid alcoholic drinks
- use caution when driving a motor vehicle or operating machinery

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- pain, nasal decongestion, or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back, or occurs with fever, rash, or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Drug Facts (continued)

Directions

- do not take more than directed (see Overdose warning)
- do not take more than 4 doses in any 24-hour period
- measure only with dosing cup provided
- do not use dosing cup with other products
- dose as follows or as directed by a doctor
- adults and children 12 years of age and over: 20 mL in dosing cup provided every 4 hours
- children under 12 years of age: do not use

Other information

- each 20 mL contains: sodium 16 mg
- store at 20-25°C (68-77°F)
- do not refrigerate

Inactive ingredients ammonium glycyrrhizate, anhydrous citric acid, D&C yellow no. 10, edetate disodium, FD&C blue no. 1, flavors, glycerin (soy), propylene glycol, sodium benzoate, sorbitol, sucralose, water, xanthan gum

Questions?

1-866-MUCINEX (1-866-682-4639)

You may also report side effects to this phone number.



HEALTH • HYGIENE • HOME

Patents: www.rb.com/patents

Tamper evident: Do not use if carton is damaged or if printed seal on blister is broken or missing.

MAXIMUM STRENGTH

Mucinex[®]
NIGHTSHIFT

SEVERE COLD & FLU

FAST RELEASE. POWERFUL SYMPTOM RELIEF!

MAXIMUM STRENGTH

NDC 72854-234-20

Mucinex[®]
NIGHTSHIFT



SEVERE COLD & FLU

Acetaminophen – Pain Reliever/Fever Reducer

Dextromethorphan HBr – Cough Suppressant

Phenylephrine HCl – Nasal Decongestant

Triprolidine HCl – Antihistamine



ACTUAL SIZE



**FAST
RELEASE**

20 CAPLETS

FOR AGES 12+

LOT:

EXP.:

MADE IN:

3176311



Questions?

1-866-MUCINEX (1-866-682-4639)

You may also report side effects to
this phone number.



HEALTH • HYGIENE • HOME

www.mucinex.com

Patents: www.rfb.com/patents

Dist. by: RB Health (US)
Parsippany, NJ 07054-0224

©2021 RB Health

3176311 011921



Drug Facts

Active ingredients (in each caplet)

Acetaminophen 325 mg.....Pain reliever/fever reducer
Dextromethorphan HBr 10 mg.....Cough suppressant
Phenylephrine HCl 5 mg.....Nasal decongestant
Tripolidine HCl 1.25 mg.....Antihistamine

Purposes

Uses

- temporarily relieves these common cold and flu symptoms:
 - cough
 - nasal congestion
 - minor aches and pains
 - sore throat
 - headache
 - sinus congestion and pressure
 - runny nose
 - sneezing
 - itching of the nose or throat
 - itchy, watery eyes due to hay fever
- temporarily reduces fever
- controls cough to help you get to sleep

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 4,000 mg in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks daily while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- liver disease
- heart disease
- diabetes
- high blood pressure
- thyroid disease
- glaucoma
- trouble urinating due to an enlarged prostate gland

Drug Facts (continued)

- a breathing problem such as emphysema or chronic bronchitis
- persistent or chronic cough such as occurs with smoking, asthma, or emphysema
- cough that occurs with too much phlegm (mucus)

Ask a doctor or pharmacist before use if you are

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers

When using this product

- do not use more than directed
- excitability may occur, especially in children
- marked drowsiness may occur
- alcohol, sedatives, and tranquilizers may increase drowsiness
- avoid alcoholic drinks
- use caution when driving a motor vehicle or operating machinery

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back, or occurs with fever, rash, or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use. **Keep out of reach of children.**

Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- do not take more than directed (see Overdose warning)
- do not take more than 8 caplets in any 24-hour period
- adults and children 12 years of age and over: take 2 caplets every 4 hours
- children under 12 years of age: do not use

Other information

- store at 20-25°C (68-77°F)

Inactive ingredients croscarmellose sodium, crospovidone, ferric oxide, hydroxypropyl cellulose, mica, microcrystalline cellulose, polyvinyl alcohol, polyvinyl alcohol polyethylene glycol copolymer, povidone, silicon dioxide, stearic acid, talc, titanium dioxide

Mucinex[®]
NIGHT SHIFT
MAXIMUM STRENGTH

SEVERE COLD & FLU

Maximum Strength per 4-hour dose
*Helps to relieve these symptoms at night
Do not take more than a total of 8 caplets in any 24-hour period.
Take only as directed.
Keep carton for full information.

PARENTS:
Learn about teen medicine abuse
www.StopMedicineAbuse.org



Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening ■ blisters ■ rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- liver disease ■ heart disease
- diabetes ■ high blood pressure
- thyroid disease ■ glaucoma
- trouble urinating due to an enlarged prostate gland
- a breathing problem such as emphysema or chronic bronchitis
- persistent or chronic cough such as occurs with smoking, asthma, or emphysema
- cough that occurs with too much phlegm (mucus)

PEEL HERE

Ask a doctor or pharmacist before use if you are

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers

When using this product

- do not use more than directed
- excitability may occur, especially in children
- marked drowsiness may occur
- alcohol, sedatives, and tranquilizers may increase drowsiness
- avoid alcoholic drinks
- use caution when driving a motor vehicle or operating machinery

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back, or occurs with fever, rash, or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Drug Facts (continued)

Directions

- do not take more than directed (see Overdose warning)
- do not take more than 4 doses in any 24-hour period
- measure only with dosing cup provided
- do not use dosing cup with other products
- dose as follows or as directed by a doctor:
- adults and children 12 years of age and older: 20 mL in dosing cup provided every 4 hours
- children under 12 years of age: do not use

Other information

- each 20 mL contains: sodium 16 mg
- store at 20-25°C (68-77°F)
- do not refrigerate

Inactive ingredients

ammonium glycyrrhizate, anhydrous citric acid, ascorbic acid, edetate disodium, FD&C blue no. 1, FD&C red no. 40, flavors, glycerin (soy), propylene glycol, sodium benzoate, sorbitol, sucralose, triacetin, triethyl citrate, water, xanthan gum

Questions?

1-866-MUCINEX (1-866-682-4639)
You may also report side effects to this phone number.



HEALTH • HYGIENE • HOME

Patents:
www.rb.com/patents
Please visit our website
www.mucinex.com



NDC 63824-270-66



**SINUS
CLEAR & COOL**

Acetaminophen – Pain Reliever/Fever Reducer
Dextromethorphan HBr – Cough Suppressant
Phenylephrine HCl – Nasal Decongestant
Triprolidine HCl – Antihistamine



**NIGHTTIME
RELIEF FOR A BETTER
MORNING**



✓ COUGH ✓ FEVER ✓ SORE THROAT
✓ RUNNY NOSE ✓ SNEEZING ✓ NASAL CONGESTION

3094737
6 FL OZ (180 mL)
FOR AGES 12+
031419



Drug Facts

Active ingredients (in each 20 mL)	Purposes
Acetaminophen 650 mg...Pain reliever/fever reducer Dextromethorphan HBr 20 mg.....Cough suppressant Phenylephrine HCl 10 mg.....Nasal decongestant Triprolidine HCl 2.5 mg.....Antihistamine	

Uses

- temporarily relieves these common cold and flu symptoms:
 - cough
 - minor aches and pains
 - headache
 - sinus congestion and pressure
 - runny nose
 - sneezing
 - itching of the nose or throat
 - itchy, watery eyes due to hay fever
- temporarily reduces fever
- controls cough to help you get to sleep

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 4000 mg in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks daily while using this product

Drug Facts (continued)

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- liver disease
- diabetes
- thyroid disease
- trouble urinating due to an enlarged prostate gland
- a breathing problem such as emphysema or chronic bronchitis
- persistent or chronic cough such as occurs with smoking, asthma, or emphysema
- cough that occurs with too much phlegm (mucus)
- heart disease
- high blood pressure
- glaucoma

PEEL HERE

Drug Facts (continued)

Ask a doctor or pharmacist before use if you are

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers

When using this product

- do not use more than directed
- excitability may occur, especially in children
- marked drowsiness may occur
- alcohol, sedatives, and tranquilizers may increase drowsiness
- avoid alcoholic drinks
- use caution when driving a motor vehicle or operating machinery

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back, or occurs with fever, rash, or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Drug Facts (continued)

Directions

- do not take more than directed (see Overdose warning)
- do not take more than 4 doses in any 24-hour period
- measure only with dosing cup provided

**PEEL CORNER TO READ COMPLETE
DRUG FACTS AND INFORMATION**

Maximum Strength per 4-hour dose
Tamper evident: Do not use if neckband
on bottle cap is broken or missing.

PARENTS:



■ do not use dosing cup with other products
■ dose as follows or as directed by a doctor
■ adults and children 12 years of age and over:
20 mL in dosing cup provided every 4 hours
■ children under 12 years of age: do not use

Other information
■ each 20 mL contains: sodium 16 mg
■ store at 20-25°C (68-77°F)
■ do not refrigerate

Inactive ingredients ammonium glycyrrhizate, anhydrous citric acid, D&C yellow no. 10, edetate disodium, FD&C blue no. 1, flavors, glycerin (soy), propylene glycol, sodium benzoate, sorbitol, sucralose, water, xanthan gum

Questions?
1-866-MUCINEX (1-866-682-4639)
You may also report side effects to this phone number.


HEALTH • HYGIENE • HOME

Patents:
www.rb.com/patents

Learn about teen medicine abuse
www.StopMedicineAbuse.org

3 63824 26267 3

Please visit our website
www.mucinex.com

Dist. by: RB Health (US)
Parsippany, NJ 07054-0224
©2019 RB Health
031519

LOT:
3094738

EXP:
MADE IN:

**POWERFUL
NIGHT TIME
RELIEF**

3094739